

Škály pre hodnotenie bolesti

Číselná škála (NRS-11)^{1,2,3}

Mierna bolesť by bola považovaná za skóre bolesti 1–3, stredná bolesť za skóre 4–7 a silná bolesť za skóre >7.⁴



Vizuálne analógová škála (VAS)^{1,5}



Tvárová škála bolesti (FPS)^{1,6}

U pacientov s určitou, aj keď obmedzenou schopnosťou komunikácie, ako sú male deti a starší pacienti, môže byť tvárová škála bolesti veľmi užitočná.⁶



1. Guidelines for the management of acute pain in emergency situations, European Society of Emergency Medicine 2020.

2. Breivik H, Borchgrevink PC, Allen SM, et al. Assessment of pain. Br J Anaesth 2008;101:17-24.

3. Downie WW, Leatham PA, Rhind VM, et al. Studies with pain rating scales. Ann Rheum Dis 1978;37:378-81.

4. Hartrick CT, Kovan JP, Shapiro S. The numeric rating scale for clinical pain measurement: a ratio measure? Pain Pract 2003;3:310-6.

5. Arthur AO, Whiteside S, Brown L, et al. Patient use of tablet computers to facilitate Emergency Department pain assessment and documentation. ISRN Emergency Medicine 2012;2012:Article ID 254530.

6. Wong-Baker FACES Foundation. Wong-Baker FACES® Pain Rating Scale. Available at <http://wongbakerfaces.org/> (Accessed January 2020).